

## Seroprevalence of *Toxoplasma gondii* Antibodies in Wild Dolphins From the Spanish Mediterranean Coast

O. Cabezón, A. R. Resendes\*, M. Domingo\*, J. A. Raga†, C. Agustí†, F. Alegre‡, J. L. Mons§, J. P. Dubey||, and S. Almería¶.

Departament de Sanitat i d'Anatomia Animals, Parasitologia i Malalties Parasitàries, Facultat de Veterinària, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain; \*Departament de Sanitat i d'Anatomia Animals, Histologia i Anatomia Patològica, Facultat de Veterinària, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain; †Departamento de Zoología e Instituto Cavanilles de Biodiversidad y Biología Evolutiva, Universidad de Valencia, P.O. Box 22085, E-46071 Valencia, Spain; ‡Centre de Recuperació d' Animals Marins de Catalunya (CRAMC), 08330 Premià de Mar, Barcelona, Spain; §Aula del Mar de Málaga, Avda. Manuel Agustín Heredia no. 35 29001 Málaga, Spain; ||Animal Parasitic Diseases Laboratory, Building 1001, Animal and Natural Resources Institute, Agricultural Research Service, United States Department of Agriculture, Beltsville, Maryland 20705-2350; ¶To whom correspondence should be addressed. e-mail: sonia.almeria@uab.es

**ABSTRACT:** Although *Toxoplasma gondii* infection has been found occasionally in cetaceans, little is known of the prevalence of antibodies to *T. gondii* in wild dolphins. Antibodies to *T. gondii* were determined in serum samples from 58 dolphins stranded in the Spanish Mediterranean coast. Modified agglutination test was used to determine *T. gondii* antibodies, and a titer of 1:25 was considered indicative of *T. gondii* infection. Antibodies to *T. gondii* were found in 4 of 36 striped dolphins (*Stenella coeruleoalba*), in 2 of 4 common dolphins (*Delphinus delphis*), in 4 of 7 bottlenose dolphins (*Tursiops truncatus*), and in 1 harbour porpoise (*Phocoena phocoena*). Antibodies were not found in 9 Risso's dolphins (*Grampus griseus*) and in 1 long-finned pilot whale (*Globicephala melas*) surveyed. The results indicate that *T. gondii* infection is frequent in at least 3 dolphin species from the Mediterranean Sea.

*Toxoplasma gondii* infections have been reported in many species of warm-blooded animals, including several species of marine mammals (Dubey and Beattie, 1988; Resendes et al., 2002; Dubey et al., 2003). Recently, concerns have been raised that *T. gondii* is a cause of mortality in southern sea otters (*Enhydra lutris nereis*), which are an endangered species in U.S. waters (Cole et al., 2000; Miller et al., 2002; Krueger et al., 2003). It has been suggested that sea otters become infected with *T. gondii* oocysts in the sea from freshwater coastal runoff (Miller et al., 2002). Although *T. gondii* infections have been found occasionally in Atlantic bottlenose dolphins (*Tursiops truncatus*) (Cruickshank et al., 1990; Inskeep et al., 1990; Di Guardo, Agrimi et al., 1995; Di Guardo, Corradi et al., 1995; Shulman et al., 1997; Jardine and Dubey, 2002), striped dolphin (*Stenella coeruleoalba*) (Domingo et al., 1992; Di Guardo, Agrimi et al., 1995; Di Guardo, Corradi et al., 1995), spinner dolphin (*S. longirostris*) (Migaki et al., 1990), Risso's dolphin (*Grampus griseus*) (Di Guardo, Agrimi et al., 1995; Resendes et al., 2002), and in a Tucuxi (*Sotalia fluviatilis*) (Bandoli and Oliveira, 1977), little is known of the prevalence of antibodies to *T. gondii* in wild dolphins. The present study reports the seroprevalence of *T. gondii* antibodies in the 4 most abundant wild species of Mediterranean dolphins: striped dolphin (*S. coeruleoalba*), bottlenose dolphin (*T. truncatus*), common dolphin (*Delphinus delphis*), and Risso's dolphin (*G. griseus*). Moreover, it also includes findings in 1 harbour porpoise (*Phocoena phocoena*) and 1 long-finned pilot whale (*Globicephala melas*).

All animals in this study were fresh dead or live-stranded on the Spanish Mediterranean coast. In necropsied dolphins, whole blood was collected from the heart, and in live-stranded animals, samples were collected from the ventral fluke vein. All samples were collected in commercial, sterile serum collector tubes and centrifuged. Sera were then frozen at  $-80^{\circ}\text{C}$  until tested. Data on species and sex were recorded for each animal. Samples were provided by different groups from cetacean stranding network: Institut Cavanilles (University of Valencia) from Comunitat Valenciana, Aula del Mar from Andalucía, and Centre de Recuperació d' Animals Marins de Catalunya (CRAMC) from Catalonia. The major part of the territorial area of the Spanish Mediterranean coast was covered by these 3 working groups. Sera were analyzed for *T. gondii* antibodies with the modified agglutination test (MAT) as described by Dubey and Desmonts (1987), with minor modifications. Briefly, sera were filtrated, using a sterile  $0.2\text{-}\mu\text{m}$  microfilter (Nalgene®, Rochester, New York), to eliminate erythrocytes from hemolysis and

any bacterial membranes present in the samples and diluted at 1:25, 1:50, and 1:500 dilutions. Hemolysis of sera is not a major problem for detection of *T. gondii* antibodies with MAT using a 1:25 dilution of serum and using the mouse-derived tachyzoites (Dubey et al., 2003). Filtration showed no interference with titration of the positive control serum used. Positive and negative controls were included in each test. Sera with doubtful results were reexamined.

Although the specificity and sensitivity of MAT have not been evaluated for the diagnosis of toxoplasmosis in marine mammals, it is the most evaluated and specific test for the diagnosis of toxoplasmosis in animals, particularly pigs (Dubey et al., 1995; Dubey, 1997). A titer of  $\geq 1:25$  was considered indicative of *T. gondii* infection in dolphins as has been for other species (Dubey and Beattie, 1988; Dubey et al., 2003). The prevalence of infection was statistically analyzed considering the variables species, habitat, and sex. It was not possible to determine species-specific age although length measurements of each animal were performed.

Data were analyzed using the SPSS 11.0 Statistical Program by chi-square test. The differences between variables were analyzed by Bonferroni or Tukey–Kramer tests. For multiple comparison, Dunn's test was used, and when variances were not homogenous, a nonparametric test was performed. The differences were considered statistically significant when  $P \leq 0.05$ . When the number of studied individuals in a species was less than 3, that species was not included in the statistical analysis.

Antibodies (MAT  $\geq 1:25$ ) to *T. gondii* were found in 4 of 7 *T. truncatus*, with titers of 1:25 in 2 and 1:50 in 2; in 2 of 4 *D. delphis*, with titer of 1:25 in 1 and 1:500 in the other; in 4 of 36 *S. coeruleoalba*, with titer of 1:25 in 3 and 1:500 in 1; and in 1 *P. phocoena*, with titer of 1:25. Antibodies to *T. gondii* were not found in sera of 9 *G. griseus* and the *G. melas*.

Species with coastal habits in the study area (*T. truncatus* and *D. delphis*) had significantly higher prevalence of *T. gondii* antibodies (6 of 11 animals, 54.4%) than pelagic species (*S. coeruleoalba* and *G. griseus*, 4 of 45 animals, 8.8%) ( $P = 0.003$ ). No significant difference was observed in the prevalence of infection and sex. Antibodies were found in 13.0% of 23 male dolphins and in 22.8% of 35 female dolphins. The postmortem studies in 14 of 58 dolphins excluded active toxoplasmosis.

This is the first report on seroprevalence of *T. gondii* infection in several cetacean species from the Mediterranean Sea. A high seroprevalence was found in *T. truncatus* and *D. delphis*, but antibodies were also detected in *S. coeruleoalba* and *P. phocoena* species. To our knowledge, prevalence of *T. gondii* antibodies has only been investigated in *D. leucas* and *T. truncatus* (Mikaelian et al., 2000; Dubey et al., 2003).

Clinical toxoplasmosis has been described in wide range of dolphin species (Bandoli and Oliveira, 1977; Cruickshank et al., 1990; Inskeep et al., 1990; Migaki et al., 1990; Domingo et al., 1992; Di Guardo, Agrimi et al., 1995; Di Guardo, Corradi et al., 1995; Shulman et al., 1997; Mikaelian et al., 2000; Jardine and Dubey, 2002; Resendes et al., 2002). Most of these cases were associated with an immunosuppressive state caused by a morbillivirus infection. *Toxoplasma gondii* has been thought to be a sporadic infection in dolphins although the seroprevalence results from this study and from the one by Dubey et al. (2003) indicate that this infection is rather frequent at least in some species.

Although no seropositive *G. griseus* was detected in our study, clinical infection has been found in 2 Mediterranean animals of this species.

The fact that higher *T. gondii* titers were found in coastal rather than in pelagic dolphins suggests that their habitat can influence the probability of infection. Marine animals could become infected by oocysts washed into the sea via coastal freshwater runoff contaminated with cat excrement. This hypothesis has been strongly supported by studies of Miller et al. (2002) in sea otters. *Toxoplasma gondii* oocysts can survive in seawater for at least 6 mo (Lindsay et al., 2003).

*Toxoplasma gondii* infection in dolphins is intriguing because they drink little water and feed on fish, squid, or other cold-blooded sea animals. Dolphins could become infected by the ingestion of water contaminated with oocysts, aquatic birds, or infected meat thrown into the sea. The role of marine invertebrates in the life cycle of *T. gondii* is unknown. *Toxoplasma gondii* is not known to parasitize any cold-blooded animal; however, molluscs can filter large quantities of water and may thus concentrate microbes from the water. Experimentally, *T. gondii* oocysts have been concentrated by molluscs and shown to remain infectious to mice (Lindsay et al., 2001; Arkush et al., 2003).

The present study indicated that *T. gondii* infection is frequent, even if it does not necessarily cause disease in western Mediterranean dolphins. Furthermore, at least 4 different Mediterranean species of dolphins can be infected, with the parasite being particularly prevalent in the coastal dolphin species.

We thank the personal from the Spanish marine mammal stranding network that support animal rescue. We also thank Elena Obón for critical suggestions to the manuscript.

#### LITERATURE CITED

- ARKUSH, K. D., M. A. MILLER, C. M. LEUTENEGGER, I. A. GARDNER, A. E. PACKAM, A. R. HECKROTH, A. TENTER, B. C. BARR, AND P. A. CONRAD. 2003. Molecular and bioassay-based detection of *Toxoplasma gondii* oocysts uptake by mussels (*Mytilus galloprovincialis*). *International Journal for Parasitology* **33**: 1087–1097.
- BANDOLI, J. G., AND C. A. B. DE OLIVEIRA. 1977. Toxoplasmosis em *Sotalia guianensis* (van Beneden, 1863), Cetacea-Delphinidae. *Folia Medica* **75**: 459–468.
- COLE, R. A., D. S. LINDSAY, D. K. HOWE, C. L. RODERICK, J. P. DUBEY, N. J. THOMAS, AND L. A. BAETEN. 2000. Biological and molecular characterizations of *Toxoplasma gondii* strains obtained from southern sea otters (*Enhydra lutris nereis*). *Journal of Parasitology* **86**: 526–530.
- CRUICKSHANK, J. J., D. M. HAINES, N. C. PALMER, AND D. J. ST. AUBIN. 1990. Cysts of a *Toxoplasma*-like organisms in an Atlantic bottlenose dolphin. *Canadian Veterinary Journal* **31**: 213–215.
- DI GUARDO, G., U. AGRIMI, L. MORELLI, G. CARDETI, S. TERRACCIANO, AND S. KENNEDY. 1995. Post mortem investigations on cetaceans found stranded on the coasts of Italy between 1990 and 1993. *Veterinary Record* **136**: 439–442.
- , A. CORRADI, U. AGRIMI, N. ZIZZO, L. MORELLI, L. PERILLO, L. KRAMER, E. CABASSI, AND S. KENNEDY. 1995. Neuropathological lesions in cetaceans found stranded from 1991 to 1993 on the coasts of Italy. *European Journal of Veterinary Pathology* **1**: 47–51.
- DOMINGO, M., J. VISA, M. PUMAROLA, A. J. MARCO, L. FERRER, R. RABANAL, AND S. KENNEDY. 1992. Pathologic and immunocytochemical studies of morbillivirus infection in striped dolphins (*Stenella coeruleoalba*). *Veterinary Pathology* **29**: 1–10.
- DUBEY, J. P. 1997. Validation of the specificity of the modified agglutination test for toxoplasmosis in pigs. *Veterinary Parasitology* **71**: 307–310.
- , AND C. P. BEATTIE. 1988. Toxoplasmosis of animals and man. CRC Press, Boca Raton, Florida, 220 p.
- , AND G. DESMONTS. 1987. Serological responses of equids fed *Toxoplasma gondii* oocysts. *Equine Veterinary Journal* **19**: 337–339.
- , P. THULLIEZ, R. M. WEIGEL, C. D. ANDREWS, P. LIND, AND E. C. POWELL. 1995. Sensitivity and specificity of various serologic tests for detection of *Toxoplasma gondii* infection in naturally infected sows. *American Journal of Veterinary Research* **56**: 1030–1036.
- , R. ZARNKE, N. J. THOMAS, S. K. WONG, W. VAN BONN, J. W. DAVIS, R. EWING, M. MENSE, O. C. H. KWOK, K. B. BECKMEN, S. ROMAND, AND P. THULLIEZ. 2003. *Toxoplasma gondii*, *Neospora caninum*, *Sarcocystis neurona*, and *Sarcocystis canis*-like infections in marine mammals. *Veterinary Parasitology* **116**: 275–296.
- INSKEEP, W., C. H. GARDINER, R. K. HARRIS, J. P. DUBEY, AND R. T. GOLDSTEIN. 1990. Toxoplasmosis in Atlantic bottle-nosed dolphin (*Tursiops truncatus*). *Journal of Wildlife Diseases* **26**: 377–382.
- JARDINE, J. E., AND J. P. DUBEY. 2002. Congenital toxoplasmosis in an Indo-Pacific bottlenose dolphin (*Tursiops aduncus*). *Journal of Parasitology* **88**: 197–199.
- KRUEDER, C., M. A. MILLER, D. A. JESSUP, J. A. LOWENSTEINE, M. D. HARRIS, J. A. AMES, T. E. CARPENTER, P. A. CONRAD, AND A. K. MAZET. 2003. Patterns of mortality in southern sea otters (*Enhydra lutris nereis*) from 1998–2001. *Journal of Wildlife Diseases* **39**: 495–509.
- LINDSAY, D. S., M. V. COLLINS, S. M. MITCHELL, R. COLE, G. J. FLICK, C. N. WETCH, A. LINDQUIST, AND J. P. DUBEY. 2003. Sporulation and survival of *Toxoplasma gondii* oocysts in sea water. *Journal of Eukaryotic Microbiology* (Suppl.) **50**: 687–688.
- , K. K. PHELPS, S. A. SMITH, G. FLICK, S. S. SUMNER, AND J. P. DUBEY. 2001. Removal of *Toxoplasma gondii* oocyst from sea water by eastern oysters (*Crassostrea virginica*). *Journal of Eukaryotic Microbiology* Suppl.: 197S–198S.
- MIGAKI, G., T. R. SAWA, AND J. P. DUBEY. 1990. Fatal disseminated toxoplasmosis in a spinner dolphin (*Stenella longirostris*). *Veterinary Pathology* **27**: 463–464.
- MIKAELIAN, I., J. BOISCLAIR, J. P. DUBEY, S. KENNEDY, AND D. MARTINEAU. 2000. Toxoplasmosis in beluga whales (*Delphinapterus leucas*) from the St. Lawrence estuary: Two case reports and a serologic survey. *Journal of Comparative Pathology* **122**: 73–76.
- MILLER, M. A., I. A. GARDNER, C. KRUEDER, D. M. PARADIES, K. R. WORCESTER, D. A. JESSUP, E. DODD, M. D. HARRIS, J. A. AMES, A. E. PACKHAM, AND P. A. CONRAD. 2002. Coastal freshwater runoff is a risk factor for *Toxoplasma gondii* infection of southern sea otters (*Enhydra lutris nereis*). *International Journal for Parasitology* **32**: 997–1006.
- RESENDES, A. R., S. ALMERÍA, J. P. DUBEY, E. OBÓN, C. JUAN-SALLÉS, E. DEGOLLADA, F. ALEGRE, O. CABEZÓN, S. PONT, AND M. DOMINGO. 2002. Disseminated toxoplasmosis in a Mediterranean pregnant Risso's dolphin (*Grampus griseus*) with transplacental fetal infection. *Journal of Parasitology* **88**: 1029–1032.
- SHULMAN, F. Y., T. P. LIPSCOMB, D. MOFFETT, A. E. KRAFFT, J. H. LICHY, M. M. TSAI, J. K. TAUBENBERGER, AND S. KENNEDY. 1997. Histologic, immunohistological, and polymerase chain reaction studies of bottlenose dolphins from the 1987–1988 United States Atlantic coast epizootic. *Veterinary Pathology* **34**: 288–295.